

File Specification for GEOS-5.7.1

Version 1.0

Release Date: 03/29/2011

Global Modeling and Assimilation Office
<http://gmao.gsfc.nasa.gov>



Global Modeling and Assimilation Office
Earth Sciences Division
NASA Goddard Space Flight Center
Greenbelt, Maryland 20771

This page intentionally left blank.

File Specification for GEOS-5.7.1 Gridded Output

Document maintained by Rob Lucchesi (GMAO, SAIC)

Permission to quote from this document should be obtained from the GMAO.

Approved by:

Michele M. Rienecker Date
Head, Global Modeling and Assimilation Office
Code 610.1, NASA GSFC

REVISION HISTORY

Version Number	Revision Date	Extent of Changes
1.0	03/29/2011	Baseline

TABLE OF CONTENTS

1. INTRODUCTION	6
2. FORMAT AND FILE ORGANIZATION.....	7
2.1 DIMENSIONS	7
2.2 VARIABLES	8
2.3 GLOBAL ATTRIBUTES	9
3. INSTANTANEOUS VS TIME-AVERAGED PRODUCTS	10
4. GRID STRUCTURE	11
4.1 HORIZONTAL STRUCTURE.....	11
4.2 VERTICAL STRUCTURE	11
5. FILE NAMING CONVENTIONS	12
5.1 FILE NAMES.....	12
5.2 EARTH SCIENCE DATA TYPES (ESDT) NAME.....	14
6. SUMMARY OF GEOS-5.6 FILE COLLECTIONS.....	16
TIME-INDEPENDENT VARIABLES	16
<i>const_2d_asm_Nx</i>	17
TIME-DEPENDENT STATE VARIABLES AND DIAGNOSTICS	18
<i>inst3_3d_asm_Np</i>	19
<i>tavg3_3d_cld_Nv</i>	20
<i>tavg3_3d_mst_Nv</i>	21
<i>tavg3_3d_mst_Ne</i>	22
<i>tavg3_3d_rad_Nv</i>	23
<i>tavg3_3d_trb_Ne</i>	24
<i>tavg3_3d_tdt_Nv</i>	25
<i>tavg3_3d_udt_Nv</i>	26
<i>tavg3_3d_qdt_Nv</i>	27
<i>tavg3_3d_odt_Nv</i>	28
<i>tavg1_2d_slv_Nx</i>	29
<i>tavg1_2d_flg_Nx</i>	31
<i>tavg1_2d_rad_Nx</i>	33
<i>tavg1_2d_lnd_Nx</i>	35
<i>tavg1_2d_ocn_Nx</i>	37
<i>tavg3_2d_aer_Nx</i>	38
<i>tavg3_2d_chm_Nx</i>	42
<i>inst3_3d_aer_Nv</i>	43
<i>inst3_3d_chm_Nv</i>	44
7. METADATA	45
7.1 GES DISC METADATA.....	45
7.2 CF METADATA	46
APPENDIX A: THE IAU PROCEDURE	47
APPENDIX B: VERTICAL STRUCTURE	49
APPENDIX C: SURFACE REPRESENTATION.....	51
REFERENCES	52
WEB RESOURCES	53
ACRONYMS.....	53

1. Introduction

The GEOS-5.7.1 Atmospheric Data Assimilation System (GEOS-5 ADAS) uses an analysis developed jointly with NOAA's National Centers for Environmental Prediction (NCEP) which allows the Global Modeling and Assimilation Office (GMAO) to take advantage of the developments at NCEP and the Joint Center for Satellite Data Assimilation (JCSDA). The GEOS-5 AGCM uses the finite-volume dynamics (Lin, 2004) integrated with various physics packages (e.g., Bacmeister et al., 2006), under the Earth System Modeling Framework (ESMF) including the Catchment Land Surface Model (CLSM) (e.g., Koster et al., 2000). The GSI analysis is a three-dimensional variational (3DVar) analysis applied in grid-point space to facilitate the implementation of anisotropic, inhomogeneous covariances (e.g., Wu et al., 2002; Derber et al., 2003). The GSI implementation for GEOS-5.7.1 incorporates a set of recursive filters that produce approximately Gaussian smoothing kernels and isotropic correlation functions.

The GEOS-5 ADAS is documented in Rienecker et al. (2008). More recent updates to the model are presented in Molod et al. (2011). The GEOS-5 system actively assimilates roughly 2×10^6 observations for each analysis, including about 7.5×10^5 AIRS radiance data. The input stream is roughly twice this volume, but because of the large volume, the data are thinned commensurate with the analysis grid to reduce the computational burden. Data are also rejected from the analysis through quality control procedures designed to detect, for example, the presence of cloud.

To minimize the spurious periodic perturbations of the analysis, GEOS-5.7.1 uses the Incremental Analysis Update (IAU) technique developed by Bloom et al. (1996). More details of this procedure are given in [Appendix A](#).

The analysis is performed at a horizontal resolution of 0.3125-degree longitude by 0.25-degree latitude and at 72 levels, extending to 0.01 hPa. All products are generated at the native resolution of the horizontal grid. The majority of data products are time-averaged, but four instantaneous products are also available. Hourly data intervals are used for two-dimensional products, while 3-hourly intervals are used for three-dimensional products. These may be on the model's native 72-layer vertical grid or at 42 pressure surfaces extending to 0.1 hPa.

This document describes the gridded output files produced by the GMAO near real-time operational GEOS-5.6.x processing. Additional details about variables listed in this file specification can be found in a separate document, the GEOS-5 File Specification Variable Definition Glossary.

2. Format and File Organization

GEOS-5.7.1 files are generated with the Network Common Data Form (NetCDF-4) library, which uses Hierarchical Data Format Version 5 (HDF-5) as the underlying format. NetCDF-4 is an open-source product of UCAR/Unidata (<https://www.unidata.ucar.edu/software/netcdf/>) and HDF-5 is developed by the HDF Group (<http://www.hdfgroup.org/>). One convenient method of reading GEOS-5 files is to use the netCDF library, but the HDF-5 library can also be used directly.

Each GEOS-5 file contains a collection of geophysical quantities that we will refer to as “fields” or “variables” as well as a set of coordinate variables that contain information about the grid coordinates. While the coordinate variables are COARDS and CF compliant, the metadata associated with the data variables may not strictly meet all CF requirements.

Due to the large size of the ¼-degree GEOS-5 data files, all products are compressed with a GRIB-like method that is invisible to the user. This method degrades the precision of the data, but every effort has been made to ensure that differences between the product and the original, uncompressed data are not scientifically meaningful. Once the precision has been degraded, the files are written using the standard gzip deflation available in NetCDF-4.

2.1 Dimensions

GMAO NetCDF-4 files contain dimension variables that can be identified and interpreted by the *units* and *positive* metadata attributes, as defined in the CF metadata conventions. The *units* attribute uses standard terminology to define specific coordinate variables, e.g., latitude, while the *positive* attribute defines whether a vertical coordinate increases or decreases from the surface to the top of the atmosphere. Some 3D products are defined on model layers rather than pressure coordinates and the units attribute is set to **layer**. This is allowed under the CF conventions to be backward compatible with the older COARDS conventions.

Table 2.1-1. Dimension Variables Contained in GMAO HDF-EOS Files

Name	Description	Type	<i>units</i> attribute	<i>positive</i> attribute (3D only)
lon	longitude	double	degrees_east	n/a
lat	latitude	double	degrees_north	n/a
lev (3D only)	pressure or layer index	double	hPa or layer	down

time	minutes since reference date & time	int	Minutes	n/a
------	-------------------------------------	-----	---------	-----

2.2 Variables

GMAO NetCDF-4 files are written using the NetCDF classic model. Arrays of scientific data are stored as variables of type **float** that contain various attributes such as *units*, *long_name*, *standard_name*, *_FillValue*, and others. Please note that we do not guarantee that the value in the *standard_name* attribute will conform to the CF metadata conventions. You can quickly list the variables as well as the complete structure of the file by using common utilities such as *ncdump* or *h5dump*. The utilities are distributed with the NetCDF and HDF distributions.

Table 2.2-1 Metadata attributes associated with each variable.

Name	Type	Description
_FillValue	float	Floating-point value used to identify missing data. Will normally be set to 1e15. Required by CF.
missing_value	float	Same as _FillValue. Included for backward compatibility.
valid_range	float32, array(2)	This attribute defines the valid range of the variable. The first element is the smallest valid value and the second element is the largest valid value. Required by CF, but this attribute is not loaded with useful data.
long_name	String	An ad hoc description of the variable as required by COARDS . It approximates the standard names as defined in an early version of CF conventions. (See References). The <i>Description</i> column from the tables of Section 6 is based on this name.
standard_name	String	Same as long_name.
Units	String	The units of the variable. Must be a string that can be recognized by UNIDATA's Udunits package.

Name	Type	Description
scale_factor	float32	If variable is packed as 16-bit integers, this is the scale_factor for expanding to floating-point. Currently we do not plan to pack data, thus value will be 1.0
add_offset	float32	If variable is packed as 16-bit integers, this is the offset for expanding to floating-point. Currently, we do not plan to pack data, thus value will be 0.0.

2.3 Global Attributes

In addition to scientific variables and dimension scales, global metadata is also stored in GMAO NetCDF-4 files. These metadata attributes are largely defined by the CF/COARDS conventions.

Table 2.3-1 Global metadata attributes associated with each variable,.

Name	Type	Description
Conventions	character	Identification of the file convention used, currently “CF”
Title	character	Experiment identification.
History	character	Processing history.
Institution	character	“NASA Global Modeling and Assimilation Office”
Source	character	CVS tag of this release. CVS tags are used internally by the GMAO to designate versions of the system.
References	character	GMAO website address
Comment	character	Standard filename of this granule.

3. Instantaneous vs Time-averaged Products

Each file collection listed in [Section 6](#) contains either instantaneous or time-averaged products, but not both.

All instantaneous collections contain fields written every 3 hours, at *synoptic times* (00 UTC, 06 UTC, 12 UTC, and 18 UTC) and at *mid-synoptic times* (03 UTC, 09 UTC, 15 UTC, and 21 UTC).

Time-averaged collections contain either hourly or three-hourly means, but not mixtures of the two. Each time-averaged collection consists of a continuous sequence of data averaged over the indicated interval and time-stamped with the central time of the interval. For hourly data, these times are 00:30 UTC, 01:30 UTC, 02:30 UTC, etc. Only products consisting exclusively of two-dimensional (horizontal) fields are produced hourly. Three-hourly time-averaged files contain averages over time intervals centered and time stamped at 01:30 UTC, 04:30 UTC, 07:30 UTC, and so on.

Each hourly and three-hourly collection is a series of files, one timestep per file, with the date and time as part of the filename.

4. Grid Structure

4.1 Horizontal Structure

Fields are produced on the model's native horizontal grid, with a resolution of 5/16 degree longitude by 1/4 degree latitude.

The GEOS-5 global horizontal grid consists of **IMn=1152** points in the longitudinal direction and **JMn=721** points in the latitudinal direction. The horizontal native grid origin, associated with variables indexed ($i=1, j=1$) represents a grid point located at (180°W, 90°S). Latitude and longitude of grid points as a function of their indices (i, j) can be determined by:

$$\lambda_i = -180 + (\Delta\lambda)_n(i-1), \quad i = 1, \text{IMn}$$

$$\varphi_j = -90 + (\Delta\varphi)_n(j-1), \quad j = 1, \text{JMn}$$

Where $(\Delta\lambda)_n = 5/16^\circ$ and $(\Delta\varphi)_n = 1/4^\circ$. For example, ($i = 577, j = 361$) corresponds to a grid point at ($\lambda = 0, \varphi = 0$).

4.2 Vertical Structure

Gridded products use four different vertical configurations: horizontal-only (can be vertical averages, single level, or surface values), pressure-level, model-level, or model-edge. Horizontal-only data for a given variable appear as 3-dimensional fields (x, y, time), while pressure-level, model-level, or model-edge data appear as 4-dimensional fields (x, y, z, time). In all cases the time dimension spans multiple files, as each file (granule) contains only one time. Pressure-level data is output on the **LMp=42** pressure levels shown in Appendix B. The model layers used for GEOS-5.7.1 products are on a terrain-following hybrid sigma-p coordinate. Model-level data is output on the **LM=72** layers shown in the second table of Appendix B. The model-edge products contain fields with **LMe = LM + 1** levels representing the layer edges. The pressure at the model top is a fixed constant, **PTOP=0.01 hPa**. Pressures at model edges should be computed by summing the DELP starting at P_{TOP}. A representative pressure for the layer can then be obtained from these. In the GEOS-4 *eta* files, one could compute the pressure on the edges by using the “ak” and “bk” values and the surface pressure. In GEOS-5, the full 3-dimensional pressure variables are explicitly provided through (DELP_{ijl}) and P_{TOP}. Even though the model-level fields are on a hybrid sigma-p coordinate and their vertical location could be obtained from the “ak-bk” relationship, this may change in future GMAO systems. We thus recommend that users rely on the reported 3D pressure distribution, and not use ones computed from the “ak” and “bk”.

Note that the indexing for the GEOS-5.7.1 vertical coordinate system is top to bottom, i.e., layer 1 is the top layer of the atmosphere, while layer LM is adjacent to the earth's surface. The same is true for edge variables, with level 1 being the top of the model's atmosphere (P_{TOP}), and level LM+1 being the surface.

5. File Naming Conventions

Each GEOS-5.7.1 product file will have a complete file name identified in the metadata attribute *comment*. EOSDIS also requires nine-character abbreviated naming indices for each Earth Science Data Type (ESDT). Each GEOS-5.7.1 file collection has a unique ESDT index. The ESDT index convention is described in section 5.2.

5.1 File Names

The standard generic complete name for the assimilated GEOS-5 products will appear as follows:

DAS.config.mode.collection.sys_ver.timestamp.file_ver.nc4

A brief description of the node fields appear below:

DAS:

Identifies output as a Data Assimilation System product.

config:

GEOS-5 will run in only one operational configuration. Other configurations may be added later.

ops - Operational assimilation, nominally 12 hours behind real-time.

mode:

GEOS-5 can run in different modes of operation, but the only mode used to support EOS instrument teams is "asm".

asm - Assimilation. Uses a combination of atmospheric data analysis and model forecasting to generate a time-series of global atmospheric quantities.

collection:

All GEOS-5.7.1 data are organized into file *collections* that contain fields with common characteristics. These collections are used to make the data more accessible for specific purposes. Fields may appear in more than one collection. Collection names are of the form *freq_dims_group_HV*, where the four attributes are:

freq: time-independent (**cnst**), instantaneous (**instF**), or time-average (**tavgF**), where *F* indicates the frequency or averaging interval and can be any of the following:

1 = Hourly

3 = 3-Hourly

dims: **2d** for collections with only 2-dimensional fields or **3d** for collections with a mix of 2- and 3-dimensional fields.

group: A three-letter mnemonic for the type of fields in the collection. It is a lowercase version of the group designation used in the ESDT name, as listed in the next section.

HV: Horizontal and Vertical grid.

H can be:

N: Native ($1/4^\circ \times 5/16^\circ$) horizontal resolution

V can be:

x: horizontal-only data (surface, single level, etc.); *dims* must be **2D**

p: pressure-level data (see Appendix B for levels); *dims* must be **3D**

v: model layer centers (see Appendix B); *dims* must be **3D**

e: model layer edges (see Appendix B); *dims* must be **3D**

sys_ver:

This node defines the GEOS-5 system version that was used to generate this data.

GEOS571 – GEOSdas 5.7.1 is the current operational version of GEOS-5.

timestamp:

This node defines the date and time associated with the data in the file. It has the form *yyyymmdd_hhmm* for either instantaneous or time-averaged daily files, *yyyymm_hhmm* for monthly-mean files.

yyyy - year string (e.g. , "2002")

mm - month string (e.g., "09" for September)

dd - day of the month string

hh – hour (UTC)

mm - minute

file_ver:

This denotes the file version in the form *V##*. Under normal conditions *##* will be 01. In the event of a processing error that requires a re-processing, this number will be incremented to identify the new version of this file.

nc4:

All files are in NetCDF-4 format, thus the suffix “.nc4”.

EXAMPLE:

DAS.ops.asm.inst3_3d_asm_Np.GEOS571.20101201_0000.V01.nc4

This is an example of a GEOS-5.7.1 filename. The collection is instantaneous, three-dimensional, assimilation products, at the native horizontal resolution, interpolated to pressure levels. The file contains data valid at 00 UTC on 01 December 2010.

5.2 Earth Science Data Types (ESDT) Name

To accommodate EOSDIS toolkit requirements, all GEOS-5 files are associated with a nine-character ESDT. The ESDT, also known as the ShortName, is a short handle for users to access sets of files and takes the form:

D57TFHVGGG

where

T: Time Description:

- I** = Instantaneous
- T** = Time-averaged
- C** = Time-independent

F: Frequency

- 1** = Hourly
- 3** = 3-Hourly

H: Horizontal Resolution

- N** = Native

V: Vertical Location

- X** = Two-dimensional
- P** = Pressure
- V** = model layer center
- E** = model layer edge

GGG: Group

- ASM** = assimilated state variables (from the IAU corrector, see [Appendix A](#))
- TDT** = tendencies of temperature
- UDT** = tendencies of eastward and northward wind components
- QDT** = tendencies of specific humidity
- ODT** = tendencies of ozone
- LND** = land surface variables
- FLX** = surface turbulent fluxes and related quantities
- MST** = moist processes
- CLD** = cloud-related quantities
- RAD** = radiation
- TRB** = turbulence
- SLV** = single level
- INT** = vertical integrals

CHM = chemistry forcing
AER = aerosol diagnostics

6. Summary of GEOS-5.7.1 file collections

The GEOS-5 forward processing (FP) product is organized into the 21 collections listed below. These are described in detail in the next sections. All data is on the model's native horizontal grid, which is a regular latitude-longitude grid with a spacing of $1/4^\circ$ in latitude and $5/16^\circ$ in longitude. Horizontal arrays (1152,721) are ordered by longitude first, with the first point at the Dateline and the South Pole, with the inner index increasing eastward. All 3d collections, except the **inst3_3d_asm_Np**, are on the model's native, hybrid sigma-p vertical grid. See Appendix B for the nominal edge pressures at the top of each layer for a surface pressure of 1000 hPa. Data in the **inst3_3d_asm_Np** are interpolated to the 42 pressure levels defined in Appendix B.

Table 6-1 - List of standard collections

Name	Description
const_2d_asm_Nx	Constant fields
inst3_3d_asm_Nv	Basic assimilated fields from IAU corrector
inst3_3d_asm_Np	Basic assimilated fields from IAU corrector
tavg3_3d_cld_Nv	Upper-air cloud related diagnostics
tavg3_3d_mst_Nv	Upper-air diagnostics from moist processes at layers
tavg3_3d_mst_Ne	Upper-air diagnostics from moist processes at layer edges
tavg3_3d_trb_Ne	Upper-air diagnostics from turbulence at layer edges
tavg3_3d_rad_Nv	Upper-air diagnostics from radiation
tavg3_3d_tdt_Nv	Upper-air temperature tendencies by process
tavg3_3d_udt_Nv	Upper-air wind tendencies by process
tavg3_3d_qdt_Nv	Upper-air humidity tendencies by process
tavg3_3d_odt_Nv	Upper-air ozone tendencies by process
tavg1_2d_slv_Nx	Single-level atmospheric state variables
tavg1_2d_flx_Nx	Surface fluxes and related quantities
tavg1_2d_rad_Nx	Surface and TOA radiative fluxes
tavg1_2d_lnd_Nx	Land related surface quantities
tavg1_2d_ocn_Nx	Ocean related surface quantities
tavg3_2d_aer_Nx	2D time-averaged aerosol diagnostics
tavg3_2d_chm_Nx	2D time-averaged chemistry diagnostics
inst3_3d_aer_Nv	3D instantaneous aerosol diagnostics
inst3_3d_chm_Nv	3D instantaneous chemistry diagnostics

Time-independent Variables

These are prescribed 2-dimensional fields that do not vary during the analysis. In Version 5.7.1,

they are available as: **DAS.ops.asm.const_2d_asm_Nx.GEOS571.00000000_0000.V01.nc4** at the full resolution.

const_2d_asm_Nx

ECS short name: D57C0NXASM
 ECS long name: GEOS571 FP 2d constants
 Characteristics: time invariant
 Dimensions: longitude: 1152, latitude: 721

Variable Name	Description	Units
PHIS	Surface geopotential	$m^2 s^{-2}$
SGH	Standard deviation of topography for gravity wave drag	m
FRLAKE	Fraction of lake type in grid box	1
FRLAND	Fraction of land type in grid box	1
FRLANDICE	Fraction of land ice type in grid box	1
FROCEAN	Fraction of ocean in grid box	1
AREA	Area of grid box	m^2

Time-dependent State Variables and Diagnostics

These histories are produced from the GCM during the last pass of the forward model during the analysis cycle. For the 3D-Var scheme used in version 5.7.1, they are from the “corrector” segment of the IAU cycle.

inst3_3d_asm_Nv

ECS short name: D57I3NVASM
ECS long name: GEOS571 FP 3d assimilated state on native levels
Characteristics: Instantaneous
Dimensions: longitude: 1152, latitude: 721, native levels: 72
Times: 00:00, 03:00, 06:00, 09:00, 12:00, 15:00, 18:00, 21:00 UTC

Variable Name	Description	Units
SLP	Sea-level pressure	Pa
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
PHIS	Surface Geopotential	$\text{m}^2 \text{s}^{-2}$
H	Geopotential height	m
O3	Ozone mixing ratio	kg kg^{-1}
QV	Specific humidity	kg kg^{-1}
QL	Cloud liquid water mixing ratio	kg kg^{-1}
QI	Cloud ice mixing ratio	kg kg^{-1}
RH	Relative humidity	1
T	Air temperature	K
U	Eastward wind component	m s^{-1}
V	Northward wind component	m s^{-1}
EPV	Ertel potential vorticity	$\text{K m}^2 \text{kg}^{-1} \text{s}^{-1}$
OMEGA	Vertical pressure velocity	Pa s^{-1}

inst3_3d_asm_Np

ECS short name: D57I3NPASM
ECS long name: GEOS571 FP 3d assimilated state on pressure levels
Characteristics: Instantaneous, on pressure levels
Dimensions: longitude: 1152, latitude: 721, native levels: 42
Times: 00:00, 03:00, 06:00, 09:00, 12:00, 15:00, 18:00, 21:00 UTC

Variable Name	Description	Units
SLP	Sea-level pressure	Pa
PS	Surface pressure	Pa
PHIS	Surface Geopotential	$\text{m}^2 \text{s}^{-2}$
H	Geopotential height	m
O3	Ozone mixing ratio	kg kg^{-1}
QV	Specific humidity	kg kg^{-1}
QL	Cloud liquid water mixing ratio	kg kg^{-1}
QI	Cloud ice mixing ratio	kg kg^{-1}
RH	Relative humidity	1
T	Air temperature	K
U	Eastward wind component	m s^{-1}
V	Northward wind component	m s^{-1}
EPV	Ertel potential vorticity	$\text{K m}^2 \text{kg}^{-1} \text{s}^{-1}$
OMEGA	Vertical pressure velocity	Pa s^{-1}

tavg3_3d_cld_Nv

ECS short name: D57T3NVCLD
ECS long name: GEOS571 FP 3d cloud diagnostics
Characteristics: Time averaged
Dimensions: longitude: 1152, latitude: 721, levels: 72
Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
RH	Relative humidity	1
QLLS	Cloud liquid water mixing ratio – large-scale	kg kg ⁻¹
QILS	Cloud ice mixing ratio – large-scale	kg kg ⁻¹
QLAN	Cloud liquid water mixing ratio – anvils	kg kg ⁻¹
QIAN	Cloud ice mixing ratio – anvils	kg kg ⁻¹
QCCU	Cloud condensate mixing ratio – convective updraft	kg kg ⁻¹
CFLS	3-D Cloud fraction – large scale	1
CFAN	3-D Cloud fraction – anvils	1
CFCU	3-D Cloud fraction – convective	1
TAUCLW	3-D Cloud optical thickness of water clouds	1
TAUCLI	3-D Cloud optical thickness of ice clouds	1

tavg3_3d_mst_Nv

ECS short name: D57T3NVMST

ECS long name: GEOS571 FP 3d moist processes diagnostics

Characteristics: Time averaged

Dimensions: longitude: 1152, latitude: 721, levels: 72

Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
DQRCU	Precipitation production rate – convective	kg kg ⁻¹ s ⁻¹
DQRLSAN	Precipitation production rate - large-scale+anvil	kg kg ⁻¹ s ⁻¹
REEVAPCN	Evaporation of precipitating convective condensate	kg kg ⁻¹ s ⁻¹
REEVAPLSAN	Evaporation of precipitating LS & anvil condensate	kg kg ⁻¹ s ⁻¹

tavg3_3d_mst_Ne

ECS short name: D57T3NEMST

ECS long name: GEOS571 FP 3d moist processes diagnostics at edges

Characteristics: Time averaged

Dimensions: longitude: 1152, latitude: 721, levels: 73

Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PLE	Air pressure	Pa
CMFMC	Upward moist convective mass flux	$\text{kg m}^{-2} \text{s}^{-1}$
PFLCU	Downward flux of liquid precipitation – convective	$\text{kg m}^{-2} \text{s}^{-1}$
PFICU	Downward flux of ice precipitation – convective	$\text{kg m}^{-2} \text{s}^{-1}$
PFLLSAN	Downward flux of liquid precipitation - large-scale+anvil	$\text{kg m}^{-2} \text{s}^{-1}$
PFILSAN	Downward flux of ice precipitation - large-scale+anvil	$\text{kg m}^{-2} \text{s}^{-1}$

tavg3_3d_rad_Nv

ECS short name: D57T3NVRAD

ECS long name: GEOS571 FP 3d radiation diagnostics

Characteristics: Time averaged

Dimensions: longitude: 1152, latitude: 721, levels: 72

Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
CLOUD	Cloud fraction	1
DTDTLWR	T tendency from terrestrial radiation	K s^{-1}
DTDTLWRCLR	T tendency from terrestrial radiation (clear sky)	K s^{-1}
DTDTSWR	T tendency from solar radiation	K s^{-1}
DTDTSWRCLR	T tendency from solar radiation (clear sky)	K s^{-1}

tavg3_3d_trb_Ne

ECS short name: D57T3NETRB
ECS long name: GEOS571 FP 3d turbulence diagnostics
Characteristics: Time averaged
Dimensions: longitude: 1152, latitude: 721, levels: 73
Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PLE	Air pressure	Pa
KM	Momentum diffusivity	$\text{m}^2 \text{s}^{-1}$
KMLS	Momentum diffusivity from Louis	$\text{m}^2 \text{s}^{-1}$
KMLK	Momentum diffusivity from Lock	$\text{m}^2 \text{s}^{-1}$
KH	Heat (scalar) diffusivity	$\text{m}^2 \text{s}^{-1}$
KHLS	Heat (scalar) diffusivity from Louis	$\text{m}^2 \text{s}^{-1}$
KHLK	Heat (scalar) diffusivity from Lock	$\text{m}^2 \text{s}^{-1}$
KHRAD	Heat (scalar) diffusivity Lock radiative contribution	$\text{m}^2 \text{s}^{-1}$
KHSFC	Heat (scalar) diffusivity Lock surface contribution	$\text{m}^2 \text{s}^{-1}$
RI	Richardson Number	Nondimensional

tavg3_3d_tdt_Nv

ECS short name: D57T3NVTDT
ECS long name: GEOS571 FP 3d temperature tendencies
Characteristics: Time averaged
Dimensions: longitude: 1152, latitude: 721, levels: 72
Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
DTDTRAD	Temperature tendency from radiation	K s^{-1}
DTDTMST	Temperature tendency from moist physics	K s^{-1}
DTDTTRB	Temperature tendency from turbulence	K s^{-1}
DTDTFRI	Temperature tendency from frictional heating	K s^{-1}
DTDTGWD	Temperature tendency from gravity wave drag	K s^{-1}
DTDTTOT	Temperature tendency from physics	K s^{-1}
DTDTDYN	Temperature tendency from dynamics	K s^{-1}
DTDTANA	Temperature tendency from analysis	K s^{-1}

tavg3_3d_udt_Nv

ECS short name: D57T3NVUDT
ECS long name: GEOS571 FP 3d wind tendencies
Characteristics: Time averaged
Dimensions: longitude: 1152, latitude: 721, levels: 72
Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
DUDTMST	U-wind tendency from moist physics	m s^{-2}
DUDTTRB	U-wind tendency from turbulence	m s^{-2}
DUDTGWD	U-wind tend from gravity wave drag	m s^{-2}
DUDTDYN	U-wind tendency from dynamics	m s^{-2}
DUDTANA	U-wind tendency from analysis	m s^{-2}
DVDTMST	V-wind tendency from moist physics	m s^{-2}
DVDTTRB	V-wind tendency from turbulence	m s^{-2}
DVDTGWD	V-wind tend from gravity wave drag	m s^{-2}
DVDTDYN	V-wind tendency from dynamics	m s^{-2}
DVDTANA	V-wind tendency from analysis	m s^{-2}
U	Eastward wind component	m s^{-1}
V	Northward wind component	m s^{-1}
OMEGA	Vertical pressure velocity	Pa s^{-1}

tavg3_3d_qdt_Nv

ECS short name: D57T3NVQDT
 ECS long name: GEOS571 FP 3d moisture tendencies
 Characteristics: Time averaged
 Dimensions: longitude: 1152, latitude: 721, levels: 72
 Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
DQVDTMST	Water vapor tendency from moist physics	kg kg ⁻¹ s ⁻¹
DQVDTTRB	Water vapor tendency from turbulence	kg kg ⁻¹ s ⁻¹
DQVDTCHM	Water vapor tendency from chemistry	kg kg ⁻¹ s ⁻¹
DQVDTDYN	Water vapor tendency from dynamics	kg kg ⁻¹ s ⁻¹
DQVDTANA	Water vapor tendency from analysis	kg kg ⁻¹ s ⁻¹
DQIDTMST	Ice tendency from moist physics	kg kg ⁻¹ s ⁻¹
DQIDTTRB	Ice tendency from turbulence	kg kg ⁻¹ s ⁻¹
DQIDTDYN	Ice tendency from dynamics	kg kg ⁻¹ s ⁻¹
DQLDTMST	Liquid water tendency from moist physics	kg kg ⁻¹ s ⁻¹
DQLDTTRB	Liquid tendency from turbulence	kg kg ⁻¹ s ⁻¹
DQLDTDYN	Liquid tendency from dynamics	kg kg ⁻¹ s ⁻¹

tavg3_3d_odt_Nv

ECS short name: D57T3NVODT
ECS long name: GEOS571 FP 3d ozone tendencies
Characteristics: Time averaged
Dimensions: longitude: 1152, latitude: 721, levels: 72
Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
DOXDTMST	Ozone tendency from moist physics	$\text{kg kg}^{-1} \text{s}^{-1}$
DOXDTTRB	Ozone tendency from turbulence	$\text{kg kg}^{-1} \text{s}^{-1}$
DOXDTCHM	Ozone tendency from chemistry	$\text{mol mol}^{-1} \text{s}^{-1}$
DOXDTDYN	Ozone tendency from dynamics	$\text{kg kg}^{-1} \text{s}^{-1}$
DOXDTANA	Ozone tendency from analysis	1e-6 s^{-1}

tavg1_2d_slv_Nx

ECS short name: D57T1NXSLV
 ECS long name: GEOS571 FP 2d single level diagnostics
 Characteristics: Time averaged, single level
 Dimensions: longitude: 1152, latitude: 721
 Times: 0:30, 1:30, 2:30, 3:30, 4:30, ... UTC

Variable	Description	Units
SLP	Sea level pressure	Pa
PS	Time averaged surface pressure	Pa
U850	Eastward wind at 850 hPa	m s ⁻¹
U500	Eastward wind at 500 hPa	m s ⁻¹
U250	Eastward wind at 250 hPa	m s ⁻¹
V850	Northward wind at 850 hPa	m s ⁻¹
V500	Northward wind at 500 hPa	m s ⁻¹
V250	Northward wind at 250 hPa	m s ⁻¹
T850	Temperature at 850 hPa	K
T500	Temperature at 500 hPa	K
T250	Temperature at 250 hPa	K
Q850	Specific humidity at 850 hPa	kg kg ⁻¹
Q500	Specific humidity at 500 hPa	kg kg ⁻¹
Q250	Specific humidity at 250 hPa	kg kg ⁻¹
H1000	Height at 1000 hPa	m
H850	Height at 850 hPa	m
H500	Height at 500 hPa	m
H250	Height at 250 hPa	m
OMEGA500	Vertical pressure velocity at 500 hPa	Pa s ⁻¹
U10M	Eastward wind at 10 m above displacement height	m s ⁻¹
U2M	Eastward wind at 2 m above the displacement height	m s ⁻¹
U50M	Eastward wind at 50 m above surface	m s ⁻¹
V10M	Northward wind at 10 m above the displacement height	m s ⁻¹
V2M	Northward wind at 2 m above the displacement height	m s ⁻¹
V50M	Northward wind at 50 m above surface	m s ⁻¹
T10M	Temperature at 10 m above the displacement height	K
T2M	Temperature at 2 m above the displacement height	K

QV10M	Specific humidity at 10 m above the displacement hght	kg kg ⁻¹
QV2M	Specific humidity at 2 m above the displacement height	kg kg ⁻¹
TS	Surface skin temperature	K
DISPH	Displacement height	m
TROPPV	PV based tropopause pressure	Pa
TROPPT	T based tropopause pressure	Pa
TROPPB	Blended tropopause pressure	Pa
TROPT	Tropopause temperature	K
TROPQ	Tropopause specific humidity	kg kg ⁻¹
CLDPRS	Cloud-top pressure	Pa
CLDTMP	Cloud-top temperature	K
TQV	Total column water vapor	kg m ⁻²
TQL	Total column cloud liquid water	kg m ⁻²
TQI	Total column cloud ice	kg m ⁻²
TOX	Total column odd oxygen	kg m ⁻²

tavg1_2d_flux_Nx

ECS short name: D57T1NXFLX
 ECS long name: GEOS571 FP 2d surface flux diagnostics
 Characteristics: Time averaged, single level
 Dimensions: longitude: 1152, latitude: 721
 Times: 0:30, 1:30, 2:30, 3:30, 4:30, ... UTC

Variable Name	Description	Units
EFLUX	Latent heat flux (positive upward)	W m^{-2}
EVAP	Surface evaporation	$\text{kg m}^{-2} \text{s}^{-1}$
HFLUX	Sensible heat flux (positive upward)	W m^{-2}
TAUX	Eastward surface wind stress	N m^{-2}
TAUY	Northward surface wind stress	N m^{-2}
TAUGWX	Eastward gravity wave surface stress	N m^{-2}
TAUGWY	Northward gravity wave surface stress	N m^{-2}
PBLH	Planetary boundary layer height	m
DISPH	Displacement height	m
BSTAR	Surface buoyancy scale	m s^{-2}
USTAR	Surface velocity scale	m s^{-1}
TSTAR	Surface temperature scale	K
QSTAR	Surface humidity scale	kg kg^{-1}
RI	Surface Richardson number	1
ZOH	Roughness length, sensible heat	m
ZOM	Roughness length, momentum	m
HLML	Height of center of lowest model layer	m
TLML	Temperature of lowest model layer	K
QLML	Specific humidity of lowest model layer	kg kg^{-1}
ULML	Eastward wind of lowest model layer	m s^{-1}
VLML	Northward wind of lowest model layer	m s^{-1}
RHOA	Surface air density	kg m^{-3}
SPEED	Effective surface wind speed	m s^{-1}
CDH	Surface exchange coefficient for heat	$\text{kg m}^{-2} \text{s}^{-1}$
CDQ	Surface exchange coefficient for moisture	$\text{kg m}^{-2} \text{s}^{-1}$
CDM	Surface exchange coefficient for momentum	$\text{kg m}^{-2} \text{s}^{-1}$
CN	Surface neutral drag coefficient	1

TSH	Effective turbulence skin temperature	K
QSH	Effective turbulence skin humidity	kg kg ⁻¹
FRSEAICE	Fraction of sea-ice	1
PRECANV	Surface precipitation flux from anvils	kg m ⁻² s ⁻¹
PRECCON	Surface precipitation flux from convection	kg m ⁻² s ⁻¹
PRECLSC	Surface precipitation flux from large-scale	kg m ⁻² s ⁻¹
PRECSNO	Surface snowfall flux	kg m ⁻² s ⁻¹
PRECTOT	Total surface precipitation flux	kg m ⁻² s ⁻¹
PGENTOT	Total generation of precipitation	kg m ⁻² s ⁻¹
PREVTOT	Total re-evaporation of precipitation	kg m ⁻² s ⁻¹

tavg1_2d_rad_Nx

ECS short name: D57T1NXRAD
 ECS long name: GEOS571 FP 2d radiation diagnostics
 Characteristics: Time averaged, single level
 Dimensions: longitude: 1152, latitude: 721
 Times: 0:30, 1:30, 2:30, 3:30, 4:30, ... UTC

Variable Name	Description	Units
EMIS	Surface emissivity	1
TS	Surface skin temperature	K
ALBEDO	Surface albedo	1
ALBNIRDF	Diffuse beam NIR surface albedo	1
ALBNIRDR	Direct beam NIR surface albedo	1
ALBVISDF	Diffuse beam VIS-UV surface albedo	1
ALBVISDR	Direct beam VIS-UV surface albedo	1
LWGEM	Emitted longwave at the surface	W m ⁻²
LWGAB	Absorbed longwave at the surface	W m ⁻²
LWGABCLR	Absorbed longwave at the surface with no clouds	W m ⁻²
LWGABCLRCLN	Absorbed longwave at the surface with no clouds or aerosol	W m ⁻²
LWGNT	Net downward longwave flux at the surface	W m ⁻²
LWGNTCLR	Net downward longwave flux at the surface for cloud-free sky	W m ⁻²
LWGNTCLRCLN	Net downward longwave flux at the surface for clear sky	W m ⁻²
LWTUP	Upward longwave flux at top of atmosphere (TOA)	W m ⁻²
LWTUPCLR	Upward longwave flux at TOA assuming clear sky	W m ⁻²
LWTUPCLRCLN	Upward longwave flux at TOA assuming clear clean sky	W m ⁻²
SWTDN	TOA incident shortwave flux	W m ⁻²
SWGDN	Surface incident shortwave flux	W m ⁻²
SWGDNCLR	Surface incident shortwave flux assuming clear sky	W m ⁻²
SWGNT	Surface net downward shortwave flux	W m ⁻²
SWGNTCLR	Surface net downward shortwave flux assuming clear sky	W m ⁻²
SWGNTCLN	Surface net downward shortwave flux assuming clean sky	W m ⁻²
SWGNTCLRCLN	Surface net downward shortwave flux assuming clear clean sky	W m ⁻²
SWTNT	TOA net downward shortwave flux	W m ⁻²
SWTNTCLR	TOA net downward shortwave flux assuming clear sky	W m ⁻²

SWTNTCLN	TOA net downward shortwave flux assuming clean sky	W m^{-2}
SWTNTCLRCLN	TOA net downward shortwave flux assuming clear clean sky	W m^{-2}
TAUHG	Optical thickness of high clouds	1
TAULOW	Optical thickness of low clouds	1
TAUMID	Optical thickness of mid-level clouds	1
TAUTOT	Optical thickness of all clouds	1
CLDHGH	High-level (above 400 hPa) cloud fraction	1
CLDLOW	Low-level (1000-700 hPa) cloud fraction	1
CLDMID	Mid-level (700-400 hPa) cloud fraction	1
CLDTOT	Total cloud fraction	1

tavg1_2d_lnd_Nx

ECS short name: D57T1NXLND
 ECS long name: GEOS571 FP 2d land surface diagnostics
 Characteristics: Time averaged, single level
 Dimensions: longitude: 1152, latitude: 721
 Times: 0:30, 1:30, 2:30, 3:30, 4:30, ... UTC

Variable Name	Description	Units
GRN	Vegetation greenness fraction	1
LAI	Leaf area index	1
GWETROOT	Root zone soil wetness	1
GWETTOP	Top soil layer wetness	1
TPSNOW	Top snow layer temperature	K
TUNST	Surface temperature of unsaturated zone	K
TSAT	Surface temperature of saturated zone	K
TWLT	Surface temperature of wilted zone	K
PRECSNO	Surface snowfall	$\text{kg m}^{-2} \text{s}^{-1}$
PRECTOT	Total surface precipitation	$\text{kg m}^{-2} \text{s}^{-1}$
SNOMAS	Snow mass	kg m^{-2}
SNODP	Snow depth	m
EVPSOIL	Bare soil evaporation	W m^{-2}
EVPTRNS	Latent heat flux due to transpiration	W m^{-2}
EVPINTR	Interception loss	W m^{-2}
EVPSBLN	Sublimation	W m^{-2}
RUNOFF	Overland runoff	$\text{kg m}^{-2} \text{s}^{-1}$
BASEFLOW	Baseflow	$\text{kg m}^{-2} \text{s}^{-1}$
SMLAND	Snowmelt	$\text{kg m}^{-2} \text{s}^{-1}$
FRUNST	Fractional unsaturated area	1
FRSAT	Fractional saturated area	1
FRSNO	Fractional snow-covered area	1
FRWLT	Fractional wilting area	1
PARDF	Surface downward PAR diffuse flux	W m^{-2}
PARDR	Surface downward PAR beam flux	W m^{-2}
SHLAND	Sensible heat flux from land	W m^{-2}
LHLAND	Latent heat flux from land	W m^{-2}

EVLAND	Evaporation from land	$\text{kg m}^{-2} \text{s}^{-1}$
LWLAND	Net downward longwave flux over land	W m^{-2}
SWLAND	Net downward shortwave flux over land	W m^{-2}
GHLAND	Downward heat flux at base of top soil layer	W m^{-2}
TWLAND	Total water store in land reservoirs	kg m^{-2}
TELAND	Energy store in all land reservoirs	J m^{-2}
WCHANGE	Total land water change per unit time	$\text{kg m}^{-2} \text{s}^{-1}$
ECHANGE	Total land energy change per unit time	W m^{-2}
SPLAND	Spurious land energy source	W m^{-2}
SPWATR	Spurious land water source	$\text{kg m}^{-2} \text{s}^{-1}$
SPSNOW	Spurious snow source	$\text{kg m}^{-2} \text{s}^{-1}$

tavg1_2d_ocn_Nx

ECS short name: D57T1NXOCN
 ECS long name: GEOS571 FP 2d ocean related variables
 Characteristics: Time averaged, single level
 Dimensions: longitude: 1152, latitude: 721
 Times: 0:30, 1:30, 2:30, 3:30, 4:30, ... UTC

Variable Name	Description	Units
U10M	Eastward wind at 10 m above displacement height	m s^{-1}
V10M	Northward wind at 10 m above the displacement height	m s^{-1}
T10M	Temperature at 10 m above the displacement height	K
QV10M	Specific humidity at 10 m above the displacement height	kg kg^{-1}
SHWTR	Sensible heat flux over open water	W m^{-2}
SHICE	Sensible heat flux over ice	W m^{-2}
HLATWTR	Latent heat flux over open water	W m^{-2}
HLATICE	Latent heat flux over ice	W m^{-2}
LWNDWTR	Net downward longwave flux over open water	W m^{-2}
LWNDICE	Net downward longwave heat flux over ice	W m^{-2}
SWNDWTR	Net downward shortwave flux over open water	W m^{-2}
SWNDICE	Net downward shortwave heat flux over ice	W m^{-2}
SNOWOCN	Snowfall over ocean	$\text{kg m}^{-2} \text{s}^{-1}$
RAINOCN	Rainfall over ocean	$\text{kg m}^{-2} \text{s}^{-1}$
TAUXWTR	Eastward component of surface stress over open water	N m^{-2}
TAUYWTR	Northward component of surface stress over open water	N m^{-2}
TAUXICE	Eastward component of surface stress over ice	N m^{-2}
TAUYICE	Northward component of surface stress over ice	N m^{-2}
FRSEAICE	Fraction of ocean covered by sea ice	1

tavg3_2d_aer_Nx

ECS short name: D57T3NXAER
ECS long name: GEOS571 FP 2d aerosol diagnostics
Characteristics: Time averaged, single level
Dimensions: longitude: 1152, latitude: 721
Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
BCEM001	Black Carbon Emission Bin 001	kg m ⁻² s ⁻¹
BCEM002	Black Carbon Emission Bin 002	kg m ⁻² s ⁻¹
BCDP001	Black Carbon Dry Deposition Bin 001	kg m ⁻² s ⁻¹
BCDP002	Black Carbon Dry Deposition Bin 002	kg m ⁻² s ⁻¹
BCWT001	Black Carbon Wet Deposition Bin 001	kg m ⁻² s ⁻¹
BCWT002	Black Carbon Wet Deposition Bin 002	kg m ⁻² s ⁻¹
BCEXTTAU	Black Carbon Extinction AOT [550 nm]	1
BCSCATAU	Black Carbon Scattering AOT [550 nm]	1
DUEM001	Dust Emission Bin 001	kg m ⁻² s ⁻¹
DUEM002	Dust Emission Bin 002	kg m ⁻² s ⁻¹
DUEM003	Dust Emission Bin 003	kg m ⁻² s ⁻¹
DUEM004	Dust Emission Bin 004	kg m ⁻² s ⁻¹
DUEM005	Dust Emission Bin 005	kg m ⁻² s ⁻¹
DUSD001	Dust Sedimentation Bin 001	kg m ⁻² s ⁻¹
DUSD002	Dust Sedimentation Bin 002	kg m ⁻² s ⁻¹
DUSD003	Dust Sedimentation Bin 003	kg m ⁻² s ⁻¹
DUSD004	Dust Sedimentation Bin 004	kg m ⁻² s ⁻¹
DUSD005	Dust Sedimentation Bin 005	kg m ⁻² s ⁻¹
DUDP001	Dust Dry Deposition Bin 001	kg m ⁻² s ⁻¹
DUDP002	Dust Dry Deposition Bin 002	kg m ⁻² s ⁻¹
DUDP003	Dust Dry Deposition Bin 003	kg m ⁻² s ⁻¹
DUDP004	Dust Dry Deposition Bin 005	kg m ⁻² s ⁻¹
DUDP005	Dust Dry Deposition Bin 005	kg m ⁻² s ⁻¹
DUWT001	Dust Wet Deposition Bin 001	kg m ⁻² s ⁻¹
DUWT002	Dust Wet Deposition Bin 002	kg m ⁻² s ⁻¹

DUWT003	Dust Wet Deposition Bin 003	$\text{kg m}^{-2} \text{ s}^{-1}$
DUWT004	Dust Wet Deposition Bin 004	$\text{kg m}^{-2} \text{ s}^{-1}$
DUWT005	Dust Wet Deposition Bin 005	$\text{kg m}^{-2} \text{ s}^{-1}$
DUSMASS	Dust Surface Mass Concentration	kg m^{-3}
DUCMASS	Dust Column Mass Density	kg m^{-2}
DUEXTTAU	Dust Extinction AOT [550 nm]	1
DUSCATAU	Dust Scattering AOT [550 nm]	1
DUSMASS25	Dust Surface Mass Concentration - PM 2.5	kg m^{-3}
DUCMASS25	Dust Column Mass Density - PM 2.5	kg m^{-2}
DUEXTT25	Dust Extinction AOT [550 nm] - PM 2.5	1
DUSCAT25	Dust Scattering AOT [550 nm] - PM 2.5	1
DUAERIDX	Dust TOMS UV Aerosol Index	1
SSEM001	Sea Salt Emission Bin 001	$\text{kg m}^{-2} \text{ s}^{-1}$
SSEM002	Sea Salt Emission Bin 002	$\text{kg m}^{-2} \text{ s}^{-1}$
SSEM003	Sea Salt Emission Bin 003	$\text{kg m}^{-2} \text{ s}^{-1}$
SSEM004	Sea Salt Emission Bin 004	$\text{kg m}^{-2} \text{ s}^{-1}$
SSEM005	Sea Salt Emission Bin 005	$\text{kg m}^{-2} \text{ s}^{-1}$
SSSD001	Sea Salt Sedimentation Bin 001	$\text{kg m}^{-2} \text{ s}^{-1}$
SSSD002	Sea Salt Sedimentation Bin 002	$\text{kg m}^{-2} \text{ s}^{-1}$
SSSD003	Sea Salt Sedimentation Bin 003	$\text{kg m}^{-2} \text{ s}^{-1}$
SSSD004	Sea Salt Sedimentation Bin 004	$\text{kg m}^{-2} \text{ s}^{-1}$
SSSD005	Sea Salt Sedimentation Bin 005	$\text{kg m}^{-2} \text{ s}^{-1}$
SSDP001	Sea Salt Dry Deposition Bin 001	$\text{kg m}^{-2} \text{ s}^{-1}$
SSDP002	Sea Salt Dry Deposition Bin 002	$\text{kg m}^{-2} \text{ s}^{-1}$
SSDP003	Sea Salt Dry Deposition Bin 003	$\text{kg m}^{-2} \text{ s}^{-1}$
SSDP004	Sea Salt Dry Deposition Bin 004	$\text{kg m}^{-2} \text{ s}^{-1}$
SSDP005	Sea Salt Dry Deposition Bin 005	$\text{kg m}^{-2} \text{ s}^{-1}$
SSWT001	Sea Salt Wet Deposition Bin 001	$\text{kg m}^{-2} \text{ s}^{-1}$
SSWT002	Sea Salt Wet Deposition Bin 002	$\text{kg m}^{-2} \text{ s}^{-1}$
SSWT003	Sea Salt Wet Deposition Bin 003	$\text{kg m}^{-2} \text{ s}^{-1}$
SSWT004	Sea Salt Wet Deposition Bin 004	$\text{kg m}^{-2} \text{ s}^{-1}$
SSWT005	Sea Salt Wet Deposition Bin 005	$\text{kg m}^{-2} \text{ s}^{-1}$
SSSMASS	Sea Salt Surface Mass Concentration	kg m^{-3}
SSCMASS	Sea Salt Column Mass Density	kg m^{-2}
SSEXTTAU	Sea Salt Extinction AOT [550 nm]	1

SSSCATAU	Sea Salt Scattering AOT [550 nm]	1
SSSMASS25	Sea Salt Surface Mass Concentration - PM 2.5	kg m ⁻³
SSCMASS25	Sea Salt Column Mass Density - PM 2.5	kg m ⁻²
SSEXTT25	Sea Salt Extinction AOT [550 nm] - PM 2.5	1
SSSCAT25	Sea Salt Scattering AOT [550 nm] - PM 2.5	1
SSAERIDX	Sea Salt TOMS UV Aerosol Index	1
SUEM001	Sulfate Emission Bin 001	kg m ⁻² s ⁻¹
SUEM002	Sulfate Emission Bin 002	kg m ⁻² s ⁻¹
SUEM003	Sulfate Emission Bin 003	kg m ⁻² s ⁻¹
SUEM004	Sulfate Emission Bin 004	kg m ⁻² s ⁻¹
SUDP001	Sulfate Dry Deposition Bin 001	kg m ⁻² s ⁻¹
SUDP002	Sulfate Dry Deposition Bin 002	kg m ⁻² s ⁻¹
SUDP003	Sulfate Dry Deposition Bin 003	kg m ⁻² s ⁻¹
SUDP004	Sulfate Dry Deposition Bin 004	kg m ⁻² s ⁻¹
SUWT001	Sulfate Wet Deposition Bin 001	kg m ⁻² s ⁻¹
SUWT002	Sulfate Wet Deposition Bin 002	kg m ⁻² s ⁻¹
SUWT003	Sulfate Wet Deposition Bin 003	kg m ⁻² s ⁻¹
SUWT004	Sulfate Wet Deposition Bin 004	kg m ⁻² s ⁻¹
SO2SMASS	SO2 Surface Mass Concentration	kg m ⁻³
SO2CMASS	SO2 Column Mass Density	kg m ⁻²
SO4SMASS	SO4 Surface Mass Concentration	kg m ⁻³
SO4CMASS	SO4 Column Mass Density	kg m ⁻²
DMSSMASS	DMS Surface Mass Concentration	kg m ⁻³
DMSCMASS	DMS Column Mass Density	kg m ⁻²
SUPSO2	SO2 Prod from DMS Oxidation [column]	kg m ⁻² s ⁻¹
SUPSO4G	SO4 Prod from Gaseous SO2 Oxidation [column]	kg m ⁻² s ⁻¹
SUPSO4AQ	SO4 Prod from Aqueous SO2 Oxidation [column]	kg m ⁻² s ⁻¹
SUPSO4WT	SO4 Prod from Aqueous SO2 Oxidation (wet dep)	kg m ⁻² s ⁻¹
SUPMSA	MSA Prod from DMS Oxidation [column]	kg m ⁻² s ⁻¹
SUEXTTAU	SO4 Extinction AOT [550 nm]	1
SUSCATAU	SO4 Scattering AOT [550 nm]	1
OCDP001	Organic Carbon Dry Deposition Bin 001	kg m ⁻² s ⁻¹
OCDP002	Organic Carbon Dry Deposition Bin 002	kg m ⁻² s ⁻¹
OCWT001	Organic Carbon Wet Deposition Bin 001	kg m ⁻² s ⁻¹
OCWT002	Organic Carbon Wet Deposition Bin 002	kg m ⁻² s ⁻¹

OCSMASS	Organic Carbon Surface Mass Concentration	kg m^{-3}
OCCMASS	Organic Carbon Column Mass Density	kg m^{-2}
OCEXTTAU	Organic Carbon Extinction AOT [550 nm]	1
OCSCATAU	Organic Carbon Scattering AOT [550 nm]	1
OCEM001	Organic Carbon Emission Bin 001	$\text{kg m}^{-2} \text{s}^{-1}$
OCEM002	Organic Carbon Emission Bin 002	$\text{kg m}^{-2} \text{s}^{-1}$
DUSV	Dust Tendency Due to Convective Scavenging	$\text{kg m}^{-2} \text{s}^{-1}$
SSSV	Sea Salt Tendency Due to Convective Scavenging	$\text{kg m}^{-2} \text{s}^{-1}$
OCSV	Organic Carbon Tendency Due to Convective Scavenging	$\text{kg m}^{-2} \text{s}^{-1}$
BCSV	Black Carbon Tendency Due to Convective Scavenging	$\text{kg m}^{-2} \text{s}^{-1}$
SUSV	Sulfate Tendency Due to Convective Scavenging	$\text{kg m}^{-2} \text{s}^{-1}$

tavg3_2d_chm_Nx

ECS short name: D57T3NXCHM
ECS long name: GEOS571 FP 2d chemistry diagnostics
Characteristics: Time averaged, single level
Dimensions: longitude: 1152, latitude: 721
Times : 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name		Units
LWI	Land-Water-Ice Mask (0=land,1=water,2=ice)	1
CO2EM	CO2 Emission	$\text{kg m}^{-2} \text{s}^{-1}$
CO2CL	CO2 Column Load	kg m^{-2}
CO2SC	CO2 Surface Concentration	1e-6
COEM	CO Emission	$\text{kg m}^{-2} \text{s}^{-1}$
COPD	CO Chemical Production	$\text{kg m}^{-2} \text{s}^{-1}$
COLS	CO Chemical Loss	$\text{kg m}^{-2} \text{s}^{-1}$
COSC	CO Surface Concentration	1e-9
COCL	CO Column Burden	kg m^{-2}

inst3_3d_aer_Nv

ECS short name: D57I3NVAER
ECS long name: GEOS571 FP 3d aerosol diagnostics
Characteristics: Instantaneous
Dimensions: longitude: 1152, latitude: 721, levels 72
Times: 0:00, 3:00, 6:00, 9:00, 12:00, 15:00, 18:00, 21:00 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
LWI	Land-Water-Ice Mask (land=0,water=1,ice=2)	1
RH	Relative Humidity	1
AIRDENS	Air Density	kg m ⁻³
DU001	Dust mixing ratio in bin 1	kg kg ⁻¹
DU002	Dust mixing ratio in bin 2	kg kg ⁻¹
DU003	Dust mixing ratio in bin 3	kg kg ⁻¹
DU004	Dust mixing ratio in bin 4	kg kg ⁻¹
DU005	Dust mixing ratio in bin 5	kg kg ⁻¹
SS001	Sea Salt mixing ratio in bin 1	kg kg ⁻¹
SS002	Sea Salt mixing ratio in bin 2	kg kg ⁻¹
SS003	Sea Salt mixing ratio in bin 3	kg kg ⁻¹
SS004	Sea Salt mixing ratio in bin 4	kg kg ⁻¹
SS005	Sea Salt mixing ratio in bin 5	kg kg ⁻¹
DMS	DMS mixing ratio	kg kg ⁻¹
SO4	SO ₄ mixing ratio	kg kg ⁻¹
SO2	SO ₂ mixing ratio	kg kg ⁻¹
SO4	SO ₄ mixing ratio	kg kg ⁻¹
MSA	MSA mixing ratio	kg kg ⁻¹
BCPHOBIC	Hydrophobic black carbon mixing ratio	kg kg ⁻¹
BCPHILIC	Hydrophilic black carbon mixing ratio	kg kg ⁻¹
OCPHOBIC	Hydrophobic organic carbon mixing ratio	kg kg ⁻¹
OCPHILIC	Hydrophilic organic carbon mixing ratio	kg kg ⁻¹

inst3_3d_chm_Nv

ECS short name: D57I3NVCHM

ECS long name: GEOS571 FP 3d chemistry diagnostics

Characteristics: Instantaneous

Dimensions: longitude: 1152, latitude: 721, levels 72

Times: 1:30, 4:30, 7:30, 10:30, 13:30, 16:30, 19:30, 22:30 UTC

Variable Name	Description	Units
PS	Surface pressure	Pa
DELP	Layer pressure thickness	Pa
AIRDENS	Air Density	kg m ⁻³
CO	Carbon monoxide mixing ratio	mol mol ⁻¹
CO2	Carbon dioxide mixing ratio	mol mol ⁻¹

7. Metadata

In addition to the metadata discussed in section 2, we add metadata intended to facilitate processing by the GES DISC. In former versions of GMAO data products, this was called EOSDIS Metadata. As discussed earlier, metadata conforming to the CF conventions is also present. In addition to what is documented here, additional metadata may be present.

7.1 GES DISC Metadata

The following metadata values will be included for each data granule (file):

Name	Description
ShortName	Short identifier for each product. Also known as ESDT. (See section 5.2)
VersionID	System version. In this case, "5.7.1"
LocalVersionID	Processing version. Usually "V01".
LocalGranuleID	Filename (See section 5.1)
Format	The data format of the product files. (NetCDF-4)
RangeBeginningDate	Start date of data in the file (YYYY-MM-DD)
RangBeginningTime	Start time of data in the file (hh:mm:ssZ)
RangeEndingDate	End date of data in the file (YYYY-MM-DD)
RangeEndingTime	End time of data in the file (hh:mm:ssZ)
NorthBoundingCoordinate	Northern extent of data grid.
WestBoundingCoordinate	Western extent of data grid.
SouthBoundingCoordinate	Southern extent of data grid.

Name	Description
EastBoundingCoordinate	Eastern extent of data grid.

These metadata values will be stored as global file attributes and most of them will be encapsulated in an ODL string appended to the file as a single global metadata attribute, *CoreMetadata.0*. This metadata string will also contain the names of each variable in the file.

7.2 CF Metadata

When [GrADS](#) or [FERRET](#) are used to view GEOS-5 gridded data sets, the application will use the CF metadata embedded in the data products. These metadata include the following information:

- Space-time grid information (dimension variables)
- Variable names and descriptions
- Variable units
- "Missing" value for each variable

Grid information and units comply with the CF conventions. Most variables, but not all, will conform to CF conventions for identification by having a valid “standard_name” attribute defined.

Appendix A: The IAU procedure

The implementation of the Incremental Analysis Update (IAU, Bloom et al., 1996) used for GEOS-5.7.1 is summarized in Figure A.1. Every six hours, at the synoptic times, an analysis is performed using backgrounds at that time, three hours earlier, and three hours later, and assimilating observations during the six-hour period spanned by the three backgrounds. The analysis increments (i.e., the difference between the analysis and the corresponding synoptic background) are then divided by a time scale (in GEOS-5.7.1 we use 6 hours) to produce an “analysis tendency.” The model is then “backed-up”, restarting it from its state three hours before the analysis time, and run for six hours, adding in the time-invariant “analysis tendency” in addition to its normal physics tendencies. At that point a restart is created that will be used next time the model is backed-up, and the first background for the next analysis cycle is saved. We refer to this first 6-hour run as the “corrector” segment of the IAU. The run is then continued without an analysis tendency for another six hours, saving the other two backgrounds needed by the next analysis---one at the next synoptic time and another at the end of the six hours. We refer to this 6-hour run as the “predictor” segment of the IAU. The entire cycle is then repeated for subsequent synoptic times. Note that during each of the four daily analysis cycles the model is run for 12 hours---a 6-hour “corrector” followed by a 6-hour “predictor.”




Figure A.1

Except for the analyses themselves, all products from GEOS-5.7.1 are produced by the model during the corrector run segment. The sequence of corrector segments (follow the green line in the figure) is a continuous model run, with the extra forcing term from the analysis tendencies. The analysis tendencies do change abruptly every six hours, but state variables are continuous (within the model’s time step) solutions of the equations of motion, albeit with the extra forcing

term.

Appendix B: Vertical Structure

Pressure-level data will be output on the following 42 pressure levels:

Level	P(hPa)										
1	1000	8	825	15	600	22	250	29	30	36	2
2	975	9	800	16	550	23	200	30	20	37	1
3	950	10	775	17	500	24	150	31	10	38	0.7
4	925	11	750	18	450	25	100	32	7	39	0.5
5	900	12	725	19	400	26	70	33	5	40	0.4
6	875	13	700	20	350	27	50	34	4	41	0.3
7	850	14	650	21	300	28	40	35	3	42	0.1

Products on the native vertical grid will be output on the following levels. Pressures are nominal for a 1000 hPa surface pressure and refer to the top edge of the layer. Note that the bottom layer has a nominal thickness of 15 hPa.

Lev	P(hPa)	Lev	P(hPa)	Lev	P(hPa)	Lev	P(hPa)	Lev	P(hPa)	Lev	P(hPa)
1	0.0100	13	0.6168	25	9.2929	37	78.5123	49	450.000	61	820.000
2	0.0200	14	0.7951	26	11.2769	38	92.3657	50	487.500	62	835.000
3	0.0327	15	1.0194	27	13.6434	39	108.663	51	525.000	63	850.000
4	0.0476	16	1.3005	28	16.4571	40	127.837	52	562.500	64	865.000
5	0.0660	17	1.6508	29	19.7916	41	150.393	53	600.000	65	880.000
6	0.0893	18	2.0850	30	23.7304	42	176.930	54	637.500	66	895.000
7	0.1197	19	2.6202	31	28.3678	43	208.152	55	675.000	67	910.000
8	0.1595	20	3.2764	32	33.8100	44	244.875	56	700.000	68	925.000
9	0.2113	21	4.0766	33	40.1754	45	288.083	57	725.000	69	940.000
10	0.2785	22	5.0468	34	47.6439	46	337.500	58	750.000	70	955.000
11	0.3650	23	6.2168	35	56.3879	47	375.000	59	775.000	71	970.000
12	0.4758	24	7.6198	36	66.6034	48	412.500	60	800.000	72	985.000

Appendix C: Surface Representation

In GEOS-5 the surface below each atmospheric column consists of a set of tiles that represent various surface types. Tiles can be of four different types: Ocean, Land, Ice, Lake, as illustrated in the figure. In each grid box a single Ice tile represents those areas covered by permanent ice. Similarly a single Lake tile represents continental areas covered permanently by water. Other continental areas (non Lake or Ice) can be further subdivided into tiles that represent parts of the grid box in different hydrological catchments, defined according to the Pfafstetter (1989) system. Each of these is, in turn, divided into subtiles (not shown in figure) that represent the wilted, unsaturated, saturated, and snow-covered fractions of the tile. These fractions vary with time and are predicted by the model based on the hydrological state of the catchment and its fine-scale topographic statistics. Details of the land model, including the partitioning into subtiles, can be found in Koster et al. (2000). The Ocean tile can be divided into two subtiles that represent the ice-covered and ice-free parts of the ocean part of the atmospheric grid box. The fractional cover of these subtiles also varies with time. In GEOS-5.7.1, the fractional cover of sea ice and the ocean surface temperature are specified from Reynolds and Smith (1994).



References

1. Bacmeister, J. T., M. J. Suarez, and F. R. Robertson, 2006. Rain Re-evaporation, Boundary Layer Convection Interactions, and Pacific Rainfall Patterns in an AGCM. *J. Atmos. Sci.*, **63**, 3383-3403.
2. Bloom, S., L. Takacs, A. DaSilva, and D. Ledvina, 1996: Data assimilation using incremental analysis updates. *Mon. Wea. Rev.*, **124**, 1256-1271.
3. Collins, N., G. Theurich, C. DeLuca, M. Suarez, A. Trayanov, V. Balaji, P. Li, W. Yang, C. Hill, and A. da Silva, 2005: Design and implementation of components in the Earth System Modeling Framework. *Int. J. High Perf. Comput. Appl.*, **19**, 341-350, DOI: 10.1177/1094342005056120.
4. Derber, J. C., R. J. Purser, W.-S. Wu, R. Treadon, M. Pondeva, D. Parrish, and D. Kleist, 2003: Flow-dependent Jb in a global grid-point 3D-Var. *Proc. ECMWF annual seminar on recent developments in data assimilation for atmosphere and ocean*. Reading, UK, 8-12 Sept. 2003.
5. Koster, R. D., M. J. Suárez, A. Ducharne, M. Stieglitz, and P. Kumar, 2000: A catchment-based approach to modeling land surface processes in a GCM, Part 1, Model Structure. *J. Geophys. Res.*, **105**, 24809-24822.
6. Molod, A., L. Takacs, M.J. Suarez, J. Bacmeister, I.S. Song, A. Eichmann, Y. Chang, 2011: The GEOS-5 Atmospheric General Circulation Model: Mean Climate and Development from MERRA to Fortuna. *Technical Report Series on Global Modeling and Data Assimilation 104606*, v**28**.
7. Pfafstetter, Otto., 1989. Classification of hydrographic basins: coding methodology, unpublished manuscript, Departamento Nacional de Obras de Saneamento, August 18, 1989, Rio de Janeiro; available from J.P. Verdin, U.S. Geological Survey, EROS Data Center, Sioux Falls, South Dakota 57198 USA. See, for example: Verdin, K.L. and J.P. Verdin, 1999, A topological system for delineation and codification of the Earth's river basins," *Journal of Hydrology*, vol. 218, nos. 1-2, pp. 1-12 or <http://gis.esri.com/library/userconf/proc01/professional/papers/pap1008/p1008.htm>
8. Reynolds, R. W., and T. M. Smith, 1994: Improved global sea surface temperature analysis using optimal interpolation. *J. Climate*, **7**, 929-948.
9. Rienecker, M.M., M.J. Suarez, R. Todling, J. Bacmeister, L. Takacs, H.-C. Liu, W. Gu, M. Sienkiewicz, R.D. Koster, R. Gelaro, I. Stajner, and E. Nielsen, 2008: The GEOS-5 Data Assimilation System - Documentation of Versions 5.0.1, 5.1.0, and 5.2.0. *Technical Report Series on Global Modeling and Data Assimilation 104606*, v**27**.
10. Wu, W.-S., R.J. Purser and D.F. Parrish, 2002: Three-dimensional variational analysis with spatially inhomogeneous covariances. *Mon. Wea. Rev.*, **130**, 2905-2916.

Web Resources

GMAO web site: <http://gmao.gsfc.nasa.gov/>

GMAO Products page: <http://gmao.gsfc.nasa.gov/products/>

NetCDF information: <http://www.unidata.ucar.edu/software/netcdf/>

CF Standard Description: <http://cf-pcmdi.llnl.gov/>

The HDF Group: <http://www.hdfgroup.org/>

Acronyms

ADAS	atmospheric data assimilation system
AOT	aerosol optical thickness
CF	Climate and Forecast metadata convention
CLSM	Catchment Land Surface Model
COARDS	Cooperative Ocean/Atmosphere Research Data Service metadata convention
DMS	dimethylsulphide
ECS	EOS Core System
EOS	Earth Observing System
ESDT	Earth Science Data Type
ESMF	Earth System Modeling Framework
FP	Forward-processing
GES DISC	Goddard Earth Sciences Data and Information Services Center
GMAO	Global Modeling and Assimilation Office
GRIB	GRIdded Binary
GSI	Gridpoint Statistical Interpolation
HDF	Hierarchical Data Format
IAU	Incremental Analysis Update
JCSDA	Joint Center for Satellite Data Assimilation
MSA	methane sulphonic acid
NCEP	National Center for Environmental Prediction
NetCDF	Network Common Data Form
PAR	photosynthetically active radiation
TOA	top of atmosphere
TOMS	Total Ozone Mapping Spectrometer
UTC	Universal Time, Coordinated