
Recent Rainfall Decline in West Africa Due to Enhanced Biomass Burning and Dust Emission




Amin Dezfuli^{1,2}, Michael Bosilovich¹, Charles Ichoku^{3,4}

Overview:


- West African monsoon (WAM), the dominant climatic feature in Africa, is most active over Southern West Africa in June.
- Livelihoods of more than 200 millions of people in SWA are affected by rainfall as they rely on rainfed agriculture, fishing, hydropower, and inland waterway systems.
- Rainfall characteristics have also health implications, for example:
- Dry & dusty conditions → high risk of meningitis
- Extreme precipitation → cholera outbreaks
- Unlike dynamical features of WAM, the role of aerosols on SWA rainfall variability is not well-understood.
- We found a two-decade negative trend in SWA's rainfall that is inversely related to the combined loadings of black carbon (BC) and dust. A possible mechanism is provided.
- Improvements in seasonal predictions, using a hybrid dynamical-statistical approach.
- The results are crucial because land use change, deforestation, and rapid urbanization growth are projected to increase aerosols in the region. Therefore, SWA could experience an increase in drought intensity and frequency in future.








- African Easterly Jet/North (AEJ-N), Tropical Easterly Jet (TEJ), Low-Level Westerly (LLW), African Easterly Wave (AEW), Saharan Heat Low (SHL): <u>Jun-Sep</u>
- AEJ-S: <u>Sep-Nov</u>
- Zonal Asymmetric Pattern of Precipitation (ZAP): <u>Dec-Mar</u>
- Atlantic Meridional Mode: <u>Mar-May</u>
- Atlantic Cold Tongue/Atlantic Niño: <u>Jun-Aug</u>
- Summer (<u>Jun-Sep</u>) & Winter (<u>Dec-Mar</u>) Indian Monsoon
- Indian Ocean Dipole (IOD) Mode: Sep-Oct


Climatic Characteristics of Southern West Africa in June **West African Monsoon**













- biomass burning presents a sharp shift to begin its most-active season.
- (6) Source of Aerosols: Local or Remote?

-15 0 15 30 **─**SWA Rain **─**M2 Dust+BC


Combined loadings of dust and BC is inversely related with rainfall for both interannual variability and particularly trend.




- Both interannual variability and trend are well-captured.
- Including aerosols adds 40 and 24% improvement in capturing the trend and interannual variability.



■ Aerosols Solar radiation absorption Heating Rainfall




- 2003 (wet): strong moist monsoon, low BC & dust transport.
- 2013 (dry): strong moist monsoon, but also extreme BC.






- Using NOAA's HYSPLIT model driven by MERRA-2 data
- Back-trajectory mode initiated from SWA at 300 m
- Calculating trajectory counts from SWA to hotspots of aerosols
- Identify regions with largest contribution to the loadings of dust and BC in SWA



- BC in SWA can be reproduced with linear combination of C emission due to fires over southern savannas of Africa and Western Sahel and the corresponding transport.
- Significant trend and interannual variability exist.
- Composites of high (2008 & 2013) and low (2003 & 2007) C transport are shown.
- Similarly, dust in SWA can also be reproduced by dust transport from two different regions over the Sahara (not shown).



Dust is from MERRA-2, rainfall from TMPA, and C from Global Fire Emissions Database (GFED). The tropical rainbelt and region of maximum biomass burning have both a meridional excursion but in opposite directions. Dust has an annual cycle, but is geographically confined to the Sahara-Sahel zone.

